

Treatment Outcomes with Internet Addicts

Dr. Kimberly S. Young, Clinical Director

Center for Internet Addiction Recovery

Published in CyberPsychology & Behavior, 2007, Vol. 10, No. 5; pp. 671-679.

Abstract

Research over the last decade has identified Internet addiction as a new and often unrecognized clinical disorder that impact a user's ability to control online use to the extent that it can cause relational, occupational, and social problems. While much of the literature explores the psychological and social factors underlying Internet addiction, little if any empirical evidence exists that examines specific treatment outcomes to deal with this new client population.

Researchers have suggested using Cognitive-Behavioral Therapy (CBT) as the treatment of choice for Internet addiction and addiction recovery in general has utilized CBT as part of treatment planning. To investigate the efficacy of using CBT with Internet addicts, this study investigated 114 clients who suffered from Internet addiction and received CBT at the Center for Online Addiction. This study employed a survey research design and outcome variables such as client motivation, online time management, improved social relationships, improved sexual functioning, engagement in off-line activities and ability to abstain from problematic applications were evaluated on the third, eighth, and twelfth sessions and over a six-month follow-up. Results suggested that Caucasian, middle-aged males, with at least a four-year degree were most likely to suffer from some form of Internet addiction. Preliminary analyses indicated that the majority of clients were able to manage their presenting complaints by the eighth session and symptom management was sustained upon a six-month follow-up. As the field of Internet addiction continues to grow, such outcome data will be useful in future treatment planning with evidenced-based protocols unique to this emergent client population.

Introduction

Technology is changing the nature of problems people are having as well as how we treat them. According to the Pew Institute, surveys fielded in 2006 show that Internet penetration among adults in the U.S. has hit an all-time high. While the percentage of Americans who say they use the Internet has fluctuated slightly, their most recent survey, fielded February 15 – April 6, 2006 showed that fully 73% of respondents (about 147 million adults) are Internet users, up from 66% (about 133 million adults) in their January 2005 survey¹. As the Internet rapidly moved into homes, schools, and businesses, early studies estimated that nearly six percent of users suffered from Internet addiction². Related studies suggested that the disorder is associated with social, academic, familial, and occupational impairment^{3 4 5}. Symptoms often identified were a preoccupation with the Internet, an inability to control use, hiding or lying about the behavior, psychological withdrawal, and continued use despite consequences of the behavior.

Over the years, the concept of Internet addiction has grown in terms of its acceptance as a legitimate clinical disorder often requiring treatment⁶. Hospitals and clinics have emerged with outpatient treatment services for Internet addiction recovery such as The Illinois Institute for Addiction at Peoria Hospital or McLean Hospital, a Harvard Medical School affiliate. In some instances, addiction rehabilitation centers have admitted cases of Internet addicts into inpatient care such as Sierra Tucson in Tucson, AZ, and college campuses have started support groups to help students who are addicted to the Internet⁷. While the mental health field has gradually developed a treatment infrastructure for Internet addiction, little has been empirically studied on specific therapy modalities and their treatment outcomes with this emergent client population.

The Dynamics of Internet Addiction

One of the most common affects of Internet addiction is frequent time distortion⁸. In his studies, Greenfield found that online addicts feel a sense of displacement when online and are unable to manage central aspects of their lives due to their growing preoccupation with online use. They start to miss important deadlines at work, spend less time with their family, and slowly withdraw from their normal routines. They neglect social connections with their friends, coworkers, and with their communities, and ultimately, their lives become unmanageable because of the Internet. As the addiction grows, they become consumed with their Internet activities, preferring online games, chatting with online friends, or gambling over the Internet, gradually ignoring family and friends in exchange for solitary time in front of the computer.

Subsequently, in a study conducted by the Stanford Institute for the Quantitative Study of Society, researchers found that use of the Internet could lead to loneliness and a decline in social engagement⁹. While controversial, the researchers concluded that “the more hours people use the Internet, the less time they spend in contact with real human beings.” In a two-year longitudinal study, Kraut and his colleagues¹⁰ randomly selected families who were given computers and instruction on Internet use, and after a two-year follow-up, higher levels of depression and loneliness were reported by respondents who used the Internet as little as a few hours a week. In fact, the study reported that greater use of the Internet led to shrinking social support and happiness, and increases in depression and loneliness. In studies specific to addictive use of the Internet, Young found that social isolation was directly associated with compulsive Internet use in nearly 56% of 396 cases of online addicts she surveyed⁴. Morhan-Martin found that loneliness was correlated with excessive online use among college students¹¹ and various studies on online

relationships, specifically online affairs, found that such virtual relationships led to marital discord, separation, and possible divorce¹².

For those suffering online addictions to pornography, sex chat rooms, or having an online affair, changes in sexual behavior off-line often occurred^{13, 14}. In several cases, Internet addicts used online sex chat rooms or online pornography to replace sexual intimacy with real-life partners, and over time, completely withdrew into the computer to meet all their sexual needs.

According to some addiction theorists^{15 16}, addictions accomplish something for the person, however illusory or momentary these benefits may actually be. Because of the mental pleasure that people find in their addictions, they begin to behave more intensely about them. For example, an alcoholic is often driven to drink at moments of excessive stress or an overeater is often driven to binge on food during moments of tension. In each case, the compulsive behavior serves to reduce the underlying emotional tension and serves as a reward for future behavior. In a similar fashion, it has been suggested that Internet addicts turn to the computer to find relief from moments of painful states of mental tension and agitation present in their lives⁶. Greenfield postulated that the Internet is not as benign as we might think and has powerful mood altering capabilities, and over 29% of the Internet addicts he studied reported using the Internet to "alter their mood or escape on a regular basis"⁸. In such instances, he found that their use of the computer was less about using it as a tool and more about finding a psychological escape to cope with life's problems. Because the addiction serves a useful purpose for the addict, the sensation or attachment can grow to such proportions that life becomes unmanageable.

Applying Cognitive Behavioral Therapy

Researchers have likened Internet addiction to addictive syndromes similar to impulse-control disorders on the Axis I Scale in the DSM^{17, 18, 19, 20, 21} and utilized various forms of DSM-IV based criteria to define Internet addiction. As an impulse control disorder, Cognitive Behavioral Therapy (CBT) has been shown to be an effective treatment for compulsive disorders such as intermittent explosive disorder, pathological gambling, and trichotillomania²². CBT has also been effective in treating substance abuse, emotional disorders, and eating disorders^{23, 24}.

CBT is a familiar treatment based on the premise that thoughts determine feelings. Patients are taught to monitor their thoughts and identify those that trigger addictive feelings and actions while they learn new coping skills and ways to prevent a relapse. CBT usually requires three to months of treatment or approximately 12 weekly sessions. The early stage of therapy is behavioral, focusing on specific behaviors and situations where the impulse control disorder causes the greatest difficulty. As therapy progresses, there is more of a focus on the cognitive assumptions and distortions that have developed and the effects of these on behavior. This involves assessment of the type of distortion, problem-solving skills and coping strategies training, modeling in therapy, support groups, and keeping thought journals²⁴.

Cognitions

Addictive thinkers, for no logical reason, will feel apprehensive, when anticipating disaster¹⁶. While addicts are not the only people who worry and anticipate negative happenings, they tend to do this more often than other people. Young suggested that this type of catastrophic thinking might contribute to compulsive Internet use in providing a psychological escape mechanism to avoid real or perceived problems²⁵. Subsequent studies hypothesized that other maladaptive cognitions such as overgeneralizing or catastrophizing, negative core beliefs, and

cognitive distortions also contribute to compulsive use of the Internet^{26, 27, 28}. Young hypothesized that those who suffer from negative core beliefs may be the ones who are drawn the most to the anonymous interactive capabilities of the Internet in order to overcome these perceived inadequacies²⁵. She suggested that cognitive restructuring should be used to address underlying negative core beliefs, cognitive distortions, and rationalizations such as “Just a few more minutes won’t hurt” for effective management of the patient’s primary symptoms.

Behavior

In cases of Internet addiction, abstinence recovery models are not practical, as computers have become such a salient part of our daily lives. Therefore, clinicians have generally agreed that moderated and controlled use of the Internet is most appropriate to treat Internet addiction. Behavior therapy is the initial focus of recovery examining both computer behavior and non-computer behavior²⁹. Computer behavior deals with actual online usage with a primary goal of abstinence from problematic applications, while retaining controlled use of the computer for legitimate purposes. For example, a lawyer addicted to Internet pornography would need to learn to abstain from adult web sites, while still being able to access the Internet to conduct legal research and to email clients. Non-computer behavior focuses on helping clients develop positive lifestyle changes for life without the Internet. Life activities that do not involve the computer are evaluated and may include relationship function, social function, or occupational function.

Young suggests using a Daily Internet Log to evaluate computer behavior and establish a baseline for clinical treatment¹⁸. Once a baseline has been established, behavior therapy is used to relearn how to use the Internet to achieve specific outcomes, such as moderated online usage and more specifically abstinence from problematic online applications and controlled use for legitimate purposes. Behavior management for both computer usage and adaptive non-computer

behavior focuses on present and overt behavior¹⁸. Techniques involved may include assertion training, behavioral rehearsal, coaching, cognitive restructuring, desensitization, modeling, reinforcement, relaxation methods, self-management, or new social skills.

While many researchers have suggested treatment approaches and recovery strategies to address Internet addiction, little has been studied on actual treatment outcomes. Therefore, this study employed a survey research design to assess the utility and application of CBT in cases of Internet addiction. This study specifically examined client treatment success to reach targeted goals based upon CBT counseling. Initially, this study examined client perceptions towards the therapist and the overall therapeutic context. Problematic online applications and problems caused by compulsive use were also evaluated to understand the nature of the client population utilized. Targeted goals included the symptoms most associated with Internet addiction such as online time management, social isolation, relationship difficulties, sexual problems, and relapse. Goals were assessed over the course of 12 sessions and at six months after treatment termination.

METHODS

Participants

Participants were 214 clients seen through the Center for Online Addiction. Candidates were screened using the Internet Addiction Test (IAT)¹⁰. The IAT is a validated testing instrument that examines symptoms of Internet addiction such as a user's preoccupation with Internet use, ability to control online use, extent of hiding or lying about online use, and continued online use despite consequences of the behavior. Clients who met the criteria were included in the study and those who exhibited high-risk behaviors such as histories of psychological trauma, sexual abuse, or Axis II pathology were excluded and sent for referrals.

Procedures

The Center for Online Addiction was established in 1995 and their web site at www.netaddiction.com provides education, support, and treatment to people concerned about Internet addiction. Clients can seek treatment through traditional outpatient services utilizing individual or family counseling services. Individual adult clients requesting treatment were screened for Internet addiction using the Internet Addiction Test. Those classified for inclusion in the study completed an intake counseling form administered during the initial session that evaluated information related to compulsive use of the Internet as part of this study.

Once an appointment had been scheduled, sessions were conducted between the client and the principle investigator. Initial sessions gathered familial background, the nature of the presenting problem, its onset, and severity. Cognitive-behavioral interventions were utilized to address presenting symptoms related to computer use, specifically abstinence from problematic online applications and strategies to control online use. Counseling also focused on behavioral issues or other underlying factors contributing to online abuse such as martial discord, job burnout, problems with coworkers, or academic troubles, depending upon the unique situation of each client. Current use of the Internet was routinely evaluated and treatment outcomes were evaluated after the third, eighth, and twelfth session and upon six-month follow-up.

Materials

The “Client Outcome Questionnaire” was constructed for the purpose of this study. Clients were administered the questionnaire after the third, eighth, and twelfth online sessions and upon a six-month follow-up. It included 12-items that addressed the client’s behavior patterns

and treatment successes during the counseling process. Questions rated how effective counseling was at helping clients' achieve targeted treatment goals associated with Internet addiction recovery. Questions assessed motivation to quit abusing the Internet; ability to control online use, engagement in off-line activities, improved relationship functioning, and improved off-line sexual functioning (if applicable). Questions were rated along a five-point Likert scale (0 = Not at all; 5 = Extremely Helpful) and assessed the quality of the counseling relationship such as: "Rate your ability to develop a supportive therapist-client relationship?" "Rate the overall quality of the counseling environment?" and "Rate the overall quality of the counseling relationship?" Questions also assessed the nature of the presenting problem and related outcome goals such as: "Rate how effective online counseling was at motivating you to abstain from abusing the Internet?" "Rate your ability to control your computer use?" "Rate how your ability to function in offline relationships has improved?" "Rate your ability to abstain from sexually explicit material on the Internet?" "Rate how your ability to engage in off-line activities away from the computer has improved?" "Rate how effective online counseling was at helping you achieve sobriety from problematic online applications?"

Instrument Validation

To determine the validity and reliability of the questionnaire, mental health practitioners in the CBT therapy field evaluated the instrument as well as through a pilot test. Three therapists whom practiced some form of CBT therapy were asked to evaluate the content of the instrument and to comment on the clarity and appropriateness of the items. Before implementing the survey, a pilot test was also administered to five randomly selected college students to check the time required to finish the questionnaire, to determine if there were ambiguity and format problems,

and clarity of questionnaire items. According to the results, the researcher made the necessary modifications. Data were analyzed and percentages with frequencies were calculated for the dichotomous items and content analyses were used to evaluate qualitative data.

RESULTS

A total of 114 clients were evaluated. Demographically, 42% of clients were women and 58% of clients were men. Mean age for males and females were quite similar, 38 and 46, respectively. 84% were Caucasian, 5% were African-American, and 11% were of Asian decent, 28% held a masters degree or doctorate, 61% held a four-year bachelors degree, and 11% had earned a high school diploma.

Table 1 outlines the problematic applications reported by clients based on gender. Men were most addicted to online pornography (30%), sexual online chat rooms (10%), online gambling (10%), and online gaming (8%), respectively. Women were most addicted to online chat rooms (30%), both sexual and general, online auction houses (4%), and in a minority of cases, one woman was addicted to online gaming (2%) and one to online shopping (2%). Results suggest that chat rooms were the most problematic online application for the general population of female clients followed by online pornography, predominantly among male clients.

Insert Table 1 Here

Table 2 categorizes the problems associated with compulsive online behavior. Time management was the most reported problem with 96% citing difficulties controlling their online use and failing at repeated attempts to cut down or curb use. Relationship problems were

reported by 85% of clients citing problems such as arguments with a spouse or significant other because of the amount of time they spent at the computer. In many cases, clients lied to others about the extent of their online use or their excessive use had caused marital problems such as separation or divorce due to romantic liaisons over the Internet or online affairs. Sexual problems were reported by 75% of clients. Specifically, they indicated a decreased interest in sexual relationships with offline partners, preferring to spend time in sex chat rooms or looking at online porn than to time normally engaged in sexual activity with a spouse. Work problems were reported by 71% of clients such as reduced job productivity, missed deadlines, and in extreme cases, job loss due to their online habit. Financial problems were reported by 42% of clients that had incurred debt due to subscription fees on online porn sites, debt due to gambling losses at virtual casinos, or too much money spent shopping online or at auction houses. Physical problems such as sleep deprivation, back strain, and eyestrain were reported by 29% of clients due to long hours sitting in front of the computer. Academic problems were reported by 15% of clients that were full-time students and had developed poor study habits, missed classes, or didn't sleep enough to concentrate on school because of their online use.

Insert Table 2 Here

Table 3 presents data related to the quality of the therapist-client relationship. In examining how comfortable clients' felt working with the therapist, clients rated their ability to develop a supportive therapist-client relationship by the third session on a Mean of 3.85 ($SD = 0.94$). By the eighth session that number increased to 4.28 ($SD = 0.71$) and by the twelfth

session, the mean increased to 4.58 ($SD = 0.46$). Sessions were terminated after the twelfth session and records were kept of clients' accounts, and upon six months after termination, clients were contacted again and re-administered the Client Outcome Questionnaire. Upon six-month follow-up, clients rated the quality of the supportive therapist-client relationship on a Mean of 3.96 ($SD = 0.73$). Clients rated the quality of the counseling environment on a Mean 3.56 ($SD = .65$) by the third session, a 4.12 ($SD = 0.76$) by the eighth session, 4.36 ($SD = 0.52$) by the twelfth session, and 3.90 ($SD = 0.65$) upon six-month follow-up. In terms of the overall quality, clients rated the therapist on a Mean of 3.35 ($SD = 1.11$), 4.25 ($SD = 0.98$) by the eighth session, 4.55 ($SD = 0.75$) by the twelfth session, and 4.10 ($SD = 0.78$) upon six-month follow-up. Results suggest that clients found the counseling environment comfortable to conduct therapeutic work and their relationship with the therapist improved the longer that they were in counseling.

Insert Table 3 Here

Table 4 outlines the average outcome goal ratings reported after the third, eighth, and twelfth sessions, and upon six month follow-up. Six factors were evaluated: motivation, ability to control use, relationship function, sexual function, time spent in off-line activities, including time with family, social outings, or time at an extracurricular hobby, and overall ability to abstain from problematic online applications. Clients' motivation to stop abusing the Internet improved along a Mean of 4.22 ($SD = 1.04$) by the third session, 3.96 ($SD = 0.54$) by the eighth session, and 4.54 ($SD = 0.52$) by the twelfth session, and 4.36 ($SD = 0.63$) upon six-month follow-up. Clients' ability to control their computer use improved along a Mean of 3.95 ($SD = 1.21$), 4.06

($SD = 0.87$) by the eighth session, 4.33 ($SD = 0.58$) by the twelfth session, and 4.22 ($SD = 0.75$) upon six-month follow-up. Clients' ability to function in off-line relationships improved by a Mean of 2.95 ($SD = 0.71$), 3.66 ($SD = 0.63$) by the eighth session, 4.42 ($SD = 0.52$) by the twelfth session, and 3.99 ($SD = 0.73$) upon six month follow-up. The ability to abstain from sexually explicit online material improved along a Mean of 2.15 ($SD = 0.91$), 2.99 ($SD = 0.94$) by the eighth session, 3.26 ($SD = 1.02$) by the twelfth session, and 3.16 ($SD = 1.12$) upon six-month follow-up. The ability to engage in off-line activities away from the computer improved along a Mean of 2.67 ($SD = 0.89$) by the third session, 4.46 ($SD = 0.74$) by the eighth session, 4.66 ($SD = 0.52$) by the twelfth session, and 4.87 ($SD = 0.55$) upon six-month follow-up. The ability of CBT counseling to help clients achieve sobriety from problematic applications improved along a Mean of 3.45 ($SD = 0.61$) by the third session, 4.28 ($SD = 0.69$) by the eighth session, 4.55 ($SD = 0.48$) by the twelfth session, and 4.35 ($SD = 0.73$) upon six-month follow-up.

Results suggest that the majority of clients showed continuous improvement by the third session, effective symptom management by the eighth and twelfth sessions, and overall improved symptom maintenance upon six-month follow up. Specifically, clients were able to maintain motivation to quit abusing the Internet and improve online time management most effectively early in the counseling process. More complex issues such as rekindling off-line relationships and engaging in off-line activities improved later into the therapy process, usually by the twelfth session. Notably, improving sexual function off-line was the most difficult outcome to achieve. Many clients noted that while they were able to abstain from sex chat or online pornography, they had troubled marital relationships, five were in the process of a divorce and thus unable to regain a satisfying sexual relationship off-line with their partners.

Insert Table 4 Here

CONCLUSIONS

A total of 114 clients were evaluated on the Internet Addiction Scale. Demographically, clients tended to be male, Caucasian, and had completed at least a four-year college degree. Interestingly, clients tended to be middle-aged and were looking for help to a situation that was relatively new in their lives. The majority of clients were able to manage their presenting complaints by the eighth session. Most were able to fully manage their symptoms by the twelfth session. Clients reported that CBT counseling was effective at ameliorating the common symptoms of online addiction: motivation to quit, online time management, social isolation, sexual dysfunction, and abstinence from problematic online applications. Upon six-month follow-up, most clients were able to maintain symptom management and continued recovery.

In this study, cognitive-behavioral techniques were applied and the data suggested that clients were able to decrease thoughts and behaviors associated with compulsive Internet use. Rationalizations that led to compulsive use were reduced, behaviors surrounding computer use in general were improved, and proactive lifestyle changes to adapt to life without the Internet were primarily achieved. Clients were able to achieve relapse prevention over a six-month period.

While this study provides one of the first studies to empirically examine the efficacy of using cognitive-behavior therapy with Internet addicts, further research should continue to investigate long-term treatment outcome effects of CBT with larger client populations. As CBT is still only one treatment approach, areas for future research should also explore systematic comparisons with other treatment modalities such as psychoanalysis, gestalt, group counseling,

or in vivo counseling within an online community to determine their therapeutic impact and efficacy. Future studies should also investigate treatment differences among the various types of Internet abuse. Subtypes of Internet addiction have been identified in the literature such as Internet gambling, online gaming, and Internet pornography addictions. Studies should examine if treatment differences exist using CBT to determine if outcomes vary along each subtype.

Further, as this study relied upon self-reported data to gage changes in online behavior, psychological health status, and social functioning, the results may be biased. Because patient self-reports may be inaccurate, future studies should include that the reports be verified by relatives or friends close to the client and/or by periodic computer monitoring to ensure greater reliability of self-reported data. Finally, as the mental health field devotes more research and resources on Internet addiction recovery, future studies should evaluate how specific treatment intervention impacts long-term recovery. Traditionally, addiction treatment programs for alcoholism and drug abuse have offered patients a mix of treatment approaches. A promising new strategy involves matching patients to interventions specific to their needs. In this same manner, matching which types of Internet addiction respond best to which treatment can increase treatment effectiveness and such treatment matching is likely to increase long-term recovery.

References

- ¹ Pew Internet and American Life Project (2006). Internet Penetration and Impact. Accessed on November 17, 2006: http://www.pewinternet.org/pdfs/PIP_Internet_Impact.pdf
- ² Greenfield, D. (1999). *Internet Addiction: Disinhibition, accelerated intimacy and other theoretical considerations*. Paper presented at the 107th annual meeting of the American Psychological Association, August 22, 1999. Boston, MA.
- ³ Brenner, V. (1997). Psychology of computer use: VLVII. Parameters of Internet use, abuse and addiction: the first 90 days of the Internet usage survey. *Psychological Reports* 80: 879-882.
- ⁴ Young, K. S. (1998) Internet addiction: The emergence of a new clinical disorder. *CyberPsychology and Behavior* 1: 237-244.
- ⁵ Morahan-Martin, J., and Schumacher, P. (1999) Incidence and correlates of pathological Internet use among college students. *Computers and Human Behavior* 16:1-17.
- ⁶ Young, K. S. (2004) Internet Addiction: The consequences of a new clinical phenomena. In: *Psychology and the New Media*. K. Doyle (Ed). pp. Thousand Oaks, CA: Am. Behavioral Scientist, pp. 1-14.
- ⁷ Kandell, J. J. (1998). Internet addiction on campus: The vulnerability of college students. *Cyberpsychology and Behavior*. 1:11-17
- ⁸ Greenfield, D.N. (1999). *Virtual addiction*, Oakland, CA: New Harbinger Publications, Inc.
- ⁹ Nie, N & Erbring, L. (2000). Debating the Societal Effects of the Internet: Connecting with the World, *Public Perspective* 11: 42-43.
- ¹⁰ Krant, R., Patterson, M., Lundmark, V., Kiesler, S., Mukopadhyay, T., & Scherlis, W. (1998) Internet Paradox: A social technology that reduces social involvement and psychological well-being? *American Psychologist* 53:1017-1031.
- ¹¹ Morahan-Martin, J. (1999) The relationship between loneliness and Internet use and abuse. *Cyber Psychology and Behavior* 2: 431-440.
- ¹² Young, K, Cooper, A., Griffin-Shelley, E., Buchanan, J, & O'Mara, J. (2000). Cybersex and Infidelity Online: Implications for evaluation and treatment. *Sexual Addiction and Compulsivity* 7: 59-74.
- ¹³ Cooper, A., Putnam, D. E., Planchon, L. A. & Boies, S. C. (1999). Online sexual compulsivity: Getting tangled in the net. *Sexual Addiction & Compulsivity: The Journal of Treatment and Prevention*, 6(2): 79-104.

¹⁴ Schneider, J. (2000). Effects of cybersex addiction on the family: Results of a survey. In: *Cybersex: The Dark Side of the Force* A. Cooper (Ed.), Philadelphia: Brunner Routledge, pp. 31-58.

¹⁵ Peele, S., & Brodsky, A. (1991). *The truth about addiction and recovery: The life process program for outgrowing destructive habits*. New York, NY: Simon & Schuster.

¹⁶ Twerski, A. (1990). *Addictive thinking: Understanding self-deception*. New York, NY: HarperCollins.

¹⁷ Young K. S. (2005). An Empirical Examination of Client Attitudes towards Online Counseling. *CyberPsychology and Behavior*. 8 (2): 172-177.

¹⁸ Young, K.S. (1998) *Caught in the Net: How to recognize the signs of Internet addiction and a winning strategy for recovery*. New York, NY: John Wiley & Sons.

¹⁹ Orzack, Hecht, M. (1999). Computer Addiction: Is it Real or is it Virtual? *Harvard Mental Health Letter*. 15 (7): 8.

²⁰ Beard, K. W., & Wolf, E. M. (2001). Modification in the proposed diagnostic criteria for Internet addiction. *CyberPsychology & Behavior* 4: 377-383.

²¹ Shapira, N. A., Lessig, M. C., Goldsmith, T. D., Szabo, S. T., Lazoritz, M., Gold, M. S., & Stein, D. J. (2003). Problematic Internet use: Proposed classification and diagnostic criteria. *Depression and Anxiety*. 17: 207-216.

²² Hucker, S.J. (2004). Disorders of impulse control. *Forensic Psychology*. In W. O'Donohue and E. Levensky (Eds.). New York, NY: Academic Press.

²³ Beck, A.T. (1979). *Cognitive Therapy and the Emotional Disorders*. Plume.

²⁴ Beck, A.T., Wright F.D., Newman, C.F., Liese, B.S. (2001). *Cognitive Therapy of Substance Abuse*. The Guilford Press.

²⁵ Young, K. S. (1999) *The Evaluation and treatment of Internet addiction*. In L. VandeCreek & T.Jackson (Eds.). Innovations in Clinical Practice: A Source Book 17,19-31. Sarasota, FL: Professional Resource Press.

²⁶ Caplan, S. E. (2002). Problematic Internet use and psychosocial well-being: Development of a theory-based cognitive-behavioral measurement instrument. *Computers in Human Behavior* 18: 553-575.

²⁷ Davis, R. A. (2001). A cognitive behavioral model of pathological Internet use. *Computers in Human Behavior*. 17: 187-195

²⁸ LaRose, R., Mastro, D., & Eastin, M. S (2001). Understanding Internet Usage: A social-cognitive approach to uses and gratifications. *Social Science Computer Review* 19 (4): 395-413.

²⁹ Hall, A. S., & Parsons, J. (2001). Internet addiction: college students case study using best practices in behavior therapy. *Journal of Mental Health Counseling*. 23:312-322.

Address reprint requests to:

Kimberly S. Young
Professor of Management Sciences
School of Business
St. Bonaventure University
St. Bonaventure, NY 14778
E-mail: kyoung@sbu.edu

Table 1: Problematic Online Applications by Gender

Online Activity	Females	Males	Total
Chat (Sexual)	30% (34)	10% (11)	40% (45)
Chat (General)	4% (5)	0% (0)	4% (5)
Pornography	0% (0)	30% (34)	30% (34)
Gambling	0% (0)	10% (11)	10% (11)
Gaming	2% (2)	8% (10)	10% (12)
Auction Houses	4% (5)	0% (0)	4% (5)
Shopping	2% (2)	0% (0)	2% (2)
Total	42% (48)	58% (65)	100% (114)

Table 2: Problems Associated with Compulsive use of the Internet

PROBLEM	YES Responses	NO Responses	BLANK Responses	TOTALS
Time	96% (109)	4% (5)	0% (0)	100% (114)
Relationship	85% (97)	15% (17)	0% (0)	100% (114)
Sexual	75% (85)	21% (24)	4% (5)	100% (114)
Work	71% (81)	29% (33)	0% (0)	100% (114)
Financial	42% (48)	58% (66)	0% (0)	100% (114)
Physical	29% (33)	71% (81)	0% (0)	100% (114)
Academic	15% (17)	81% (92)	4% (5)	100% (114)

Table 3: Ratings of Therapist-Client Interaction over Three, Eight, Twelve Sessions and Six Month Follow-up

Therapy Variable	Mean Rating of Therapy Variables Measured (SD)			
	3 rd Session	8 th Session	12 th Session	Six Months
Ability to develop a supportive therapist-client relationship				
M	3.85	4.28	4.58	3.96
SD	0.94	0.71	0.46	0.73
Overall quality of the Counseling Environment				
M	3.56	4.12	4.36	3.90
SD	0.71	0.76	0.52	0.91
Quality of the Overall Counseling Relationship				
M	3.35	4.25	4.55	4.10
SD	1.11	0.98	0.75	0.78

**Table 4: Outcome Goals Assessed over
Three, Eight, Twelve Sessions and Six Month Follow-up**

Outcome Variable	Mean Rating of Outcome Variables Measured (SD)			
	3 rd Session	8 th Session	12 th Session	Six Months
Motivation				
M	4.22	3.96	4.54	4.36
SD	1.04	0.54	0.52	0.63
Time Management				
M	3.95	4.06	4.33	4.22
SD	1.21	0.87	0.58	0.75
Relationship Function				
M	2.95	3.66	4.42	3.99
SD	0.71	0.63	0.52	0.73
Sexual Function				
M	2.15	2.99	3.26	3.16
SD	0.91	0.94	1.02	1.12

Engage in Off-line Activities

M	2.67	4.46	4.66	4.87
SD	0.89	0.74	0.52	0.55

Abstinence from Problematic Applications

M	3.45	4.28	4.55	4.35
SD	0.61	0.69	0.45	0.73